

Equivalent sets of solutions of the Dirac equation with a constant electric field

A.I.Nikishov [†]

[†]*I.E.Tamm Department of Theoretical Physics, Lebedev Physical Institute,
 Leninsky Prospekt 53, 119991, Moscow, Russia
 e-mail: nikishov@lpi.ru*

Abstract

Two families of sets, nonstationary and stationary, are obtained. Each nonstationary set ψ_{p_v} consists of the solutions with the quantum number $p_v = p^0 v - p_3$. It can be obtained from the nonstationary set ψ_{p_3} with quantum number p_3 by a boost along x_3 -axis (along the direction of the electric field) with velocity $-v$. Similarly, any stationary set of solutions characterized by a quantum number $p_s = p^0 - sp_3$ can be obtained from stationary solutions with quantum number p^0 by the same boost with velocity $-s$. All these sets are equivalent and the classification (i.e. ascribing the frequency sign and in-, out- indexes) in any set is determined by the classification in ψ_{p_3} -set, where it is beyond doubt.

1 Introduction

The problems connected with Klein paradox and pair production by an external field were considered in [1-5]. The choice of the in- and out- solutions of the Klein-Gordon and Dirac equations with one-dimensional potential in [1-2] (see also [3]) disagrees with that in [4-5]. In this connection we studied in [6] the families of solutions of the Klein-Gordon equation with constant electric field and showed that our choice is correct. The existence of families of solutions and their properties are of interest by itself and here we show that the similarity of scalar and spinor cases is indeed very close.

2 Nonstationary sets

We use the metric

$$\eta_{\mu\nu} = \text{diag}(-1, 1, 1, 1), \quad (1)$$

and denote the electron charge $e' = -e = -|e|$. First we show how to obtain the nonstationary solutions characterized by the set of quantum numbers $n = (r, p_1, p_2, p_v = p^0 v - p_3)$;

$r = 1, 2$ indicates spin state. For this we take vector-potential as

$$A_\mu = -\delta_{\mu 3} E x_v, \quad x_v = t - v x_3, \quad 0 \leq v < 1. \quad (2)$$

The squared Dirac equation

$$(\Pi^2 + m^2 + g)F = 0, \quad \Pi_\mu = -i \frac{\partial}{\partial x^\mu} + e A_\mu \quad (3)$$

differs from the Klein-Gordon equation only by the presence of matrix g , which we take in the spinor representation

$$g = -ieE\alpha_3 = -ieE \begin{pmatrix} \sigma_3 & 0 \\ 0 & -\sigma_3 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (4)$$

As g is diagonal in this representation, we see that the solution F has the form

$$F = \exp[i\vartheta_v] \text{diag}(\phi_1, \phi_2, \phi_2, \phi_1), \quad (5)$$

where ϑ_v is (see eq.(18) in [6] with $e \rightarrow -e$)

$$\vartheta_v = \frac{p_v(tv - x_3)}{1 - v^2} + \frac{eEvx_v^2}{2(1 - v^2)}. \quad (6)$$

The motion along x_1, x_2 remains free and we drop the corresponding factor $\exp[ip_1x_1 + ip_2x_2]$ for brevity.

It follows from (3) and (4) that ϕ_1, ϕ_2 can be obtained from the solutions of the Klein-Gordon equation by substitutions

$$m^2 \rightarrow m^2 \mp ieE. \quad (7)$$

We denote the positive-frequency in- (out-) solution of eq. (3) as ${}_+F$ (${}_+F$) and similarly for negative-frequency solutions. Now we are in a position to find the solutions of this equation. First we note that the corresponding solutions of the Klein-Gordon equation are (see eq. (22) in [6])

$$\begin{aligned} \pm \psi_{p_v} &= e^{i\vartheta_v} D_{\pm i\kappa - \frac{1}{2}}(-e^{\mp \frac{\pi}{4}} T_v), \quad \pm \psi_{p_v} = e^{i\vartheta_v} D_{\mp i\kappa - \frac{1}{2}}(e^{\pm \frac{\pi}{4}} T_v), \\ T_v &= \sqrt{\frac{2}{eE(1 - v^2)}}(p_v + eEx_v), \quad \kappa = \frac{m^2 + p_1 + p_2}{2eE}. \end{aligned} \quad (8)$$

Applying the substitution (7), e. g., to ${}_+ \psi_{p_v} = e^{i\vartheta_v} D_{-i\kappa - \frac{1}{2}}(e^{i\frac{\pi}{4}} T_v)$, we find

$${}_+ F = e^{i\vartheta_v} \text{diag}(D_{-i\kappa - 1}(\chi_v), D_{-i\kappa}(\chi_v), D_{-i\kappa}(\chi_v), D_{-i\kappa - 1}(\chi_v)), \quad \chi_v = e^{i\frac{\pi}{4}} T_v. \quad (9)$$

The solutions of the Dirac equation are obtained as the columns of the matrix

$$(m - i\hat{\Pi})F = \begin{pmatrix} m & 0 & \Pi^0 + \Pi_3 & p_1 - ip_2 \\ 0 & m & p_1 + ip_2 & \Pi^0 - \Pi_3 \\ \Pi^0 - \Pi_3 & -p_1 + ip_2 & m & 0 \\ -p_1 - ip_2 & \Pi^0 + \Pi_3 & 0 & m \end{pmatrix} F. \quad (10)$$

Here

$$\Pi^0 = i\frac{\partial}{\partial t}, \quad \Pi_3 = -i\frac{\partial}{\partial x_3} - e'A_3 = -i\frac{\partial}{\partial x_3} - eEx_v. \quad (11)$$

It is easy to verify that

$$(\Pi^0 \pm \Pi_3)e^{i\vartheta_v}\phi(\chi_v) = -e^{i\vartheta_v - \frac{i\pi}{4}}\sqrt{\frac{2eE(1 \pm v)}{1 \mp v}}\left(\frac{\partial}{\partial \chi_v} \pm \frac{\chi_v}{2}\right)\phi(\chi_v). \quad (12)$$

We use the fourth (the first) column in matrix (10) to obtain the first (the second) spin state ${}^+\psi_{1p_v}$ (${}^+\psi_{2p_v}$). Taking into account also the first relation in

$$\left(\frac{d}{dz} - \frac{z}{2}\right)D_\nu(z) = -D_{\nu+1}(z), \quad \left(\frac{d}{dz} + \frac{z}{2}\right)D_\nu(z) = \nu D_{\nu-1}(z), \quad (13)$$

(see eqs. (8.2.15-16) in [7]), we find

$${}^+\psi_{1p_v}(x|A_3 = -Ex_v) \propto \begin{bmatrix} (p_1 - ip_2)D_{-i\kappa-1}(\chi_v) \\ e^{-\frac{i\pi}{4}}\sqrt{\frac{2eE(1-v)}{1+v}}D_{-i\kappa}(\chi_v) \\ 0 \\ mD_{-i\kappa-1}(\chi_v) \end{bmatrix} e^{i\vartheta_v}, \quad (14)$$

$${}^+\psi_{2p_v}(x|A_3 = -Ex_v) \propto \begin{bmatrix} mD_{-i\kappa-1}(\chi_v) \\ 0 \\ e^{-\frac{i\pi}{4}}\sqrt{\frac{2eE(1-v)}{1+v}}D_{-i\kappa}(\chi_v) \\ -(p_1 + p_2)D_{-i\kappa-1}(\chi_v) \end{bmatrix} e^{i\vartheta_v}, \quad (15)$$

From (14-15) for $v = 0$ we obtain the usual nonstationary solutions ψ_{p_3} with quantum number p_3 . On the other hand the solutions ψ_{p_v} can be obtained from ψ_{p_3} by a boost along x_3 with velocity $-v$. Then ψ_{p_3} is multiplied by the matrix [8]

$$\cosh \frac{\varphi}{2} - \alpha_3 \sinh \frac{\varphi}{2}, \quad \varphi = \frac{1}{2} \ln \frac{1+v}{1-v}. \quad (16)$$

In spinor representation this matrix has the form (see eq. (4) for α_3 in this representation)

$$\text{diag}(e^{-\frac{\varphi}{2}}, e^{\frac{\varphi}{2}}, e^{\frac{\varphi}{2}}, e^{-\frac{\varphi}{2}}) = \left(\frac{1-v}{1+v}\right)^{-1/4} \text{diag}\left(1, \left(\frac{1-v}{1+v}\right)^{1/2}, \left(\frac{1-v}{1+v}\right)^{1/2}, 1\right). \quad (17)$$

The transition between the standard and spinor representations is made with the help of matrix [8]

$$U = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}, \quad I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad U^2 = 1. \quad (18)$$

For $v = 1$ we get the solutions with quantum number $p^- = p^0 - p_3$. In the standard representation we find

$${}^+\psi_{rp^-}(x|A_3 = -eEx^-) = \frac{1}{2}\left[u'_r + \frac{u_r}{\pi^-}\right] \exp\left\{i\left[-\frac{p^-x^+}{2} - \frac{1}{4}eE(x^-)^2 - \kappa \ln z\right]\right\},$$

$$_-\psi_{rp^-}(x|A_3 = -eEx^-) = -\frac{1}{2}\left[u'_r + \frac{u_r}{\pi^-}\right] \exp\left\{i\left[-\frac{p^-x^+}{2} - \frac{1}{4}eE(x^-)^2 - \varkappa \ln(-z)\right]\right\}. \quad (19)$$

Here

$$z = \frac{\pi^-}{\sqrt{eE}}, \quad \pi^- = p^- + eEx^-, \quad x^\pm = t \pm x_3, \quad r = 1, 2,$$

$$u'_1 = \begin{bmatrix} 0 \\ 1 \\ 0 \\ 1 \end{bmatrix}, \quad u'_2 = \begin{bmatrix} 1 \\ 0 \\ -1 \\ 0 \end{bmatrix}, \quad u_1 = \begin{bmatrix} p_1 - ip_2 \\ m \\ p_1 - ip_2 \\ -m \end{bmatrix}, \quad u_2 = \begin{bmatrix} m \\ -p_1 - ip_2 \\ m \\ p_1 + ip_2 \end{bmatrix}. \quad (20)$$

As in the scalar case [6], we have

$$\exp[-i\varkappa \ln z] = e^{-\pi\varkappa} \exp[-i\varkappa \ln(-z)], \quad z < 0;$$

$$\exp[-i\varkappa \ln(-z)] = e^{-\pi\varkappa} \exp[-i\varkappa \ln z], \quad z > 0. \quad (21)$$

Two other solutions are

$$_+\psi_{rp^-} = \theta(\pi^-)c_{1n}^+ \psi_{rp^-}, \quad _-\psi_{rp^-} = \theta(-\pi^-)c_{1n}^- \psi_{rp^-}, \quad n = (r, p_1, p_2, p^-),$$

$$\theta(x) = \begin{cases} 1, & x > 0 \\ 0, & x < 0, \end{cases} \quad c_{1n} = i\sqrt{\frac{2\pi}{\varkappa}} \frac{\exp[-\frac{\pi\varkappa}{2}]}{\Gamma(-i\varkappa)}. \quad (22)$$

For any v the in- and out-solutions are related by Bogoliubov transformations

$$_+\psi_{rp_v} = c_{1n}^+ \psi_{rp_v} + c_{2n}^+ \psi_{rp_v},$$

$$_-\psi_{rp_v} = -c_{2n}^{*-} \psi_{rp_v} + c_{1n}^{*-} \psi_{rp_v}, \quad (23)$$

were

$$|c_{1n}|^2 + |c_{2n}|^2 = 1 \quad c_{2n} = e^{-\pi\varkappa}, \quad n = (r, p_1, p_2, p_v). \quad (23a)$$

The Bogoliubov coefficients c_{1n} , c_{2n} depend only on $\varkappa$, defined in (8).

As in the scalar case [6], the solutions with quantum number p_v can be obtained from the solutions with quantum number p^- by the integral transformation

$$\psi_{rp_v} = \int_{-\infty}^{\infty} dp^- N(p_v, p^-) \psi_{rp^-}, \quad (24)$$

$$N(p_v, p^-) = (2\pi eE(1-v))^{-\frac{1}{2}} \exp\left\{i \frac{[p^-(1+v)]^2 - 4p^-p_v(1+v) + 2p_v^2}{4eE(1-v^2)}\right\}. \quad (25)$$

Thus, using in the right-hand side of (24) the function $_+\psi_{rp^-}(x|A_3 = -Ex^-)$ from (19), we get $_+\psi_{p_v}(x|A_3 = -Ex^-)$ in the left-hand side. Changing the potential to $A_3 = -Et$, we find in the standard representation

$$_+\psi_{rp_v}(x|A_3 = -Et) = \int_{-\infty}^{\infty} dp^- N(p_v, p^-) _+\psi_{rp^-}(x|A_3 = -Et) =$$

$$\frac{\exp[-\frac{\pi\varkappa}{4}]}{2\sqrt{eE(1-v)}} [u_r D_{-i\varkappa-1}(\chi_v) + e^{-\frac{i\pi}{4}} \sqrt{\frac{2eE(1-v)}{1+v}} u'_r D_{-i\varkappa}(\chi_v)] \times$$

$$\exp\{i[\frac{\pi}{2} + \frac{\varkappa}{2} \ln \frac{1+v}{2(1-v)} + \tilde{\vartheta}_v]\}, \quad \tilde{\vartheta}_v = \vartheta_v + \frac{eEvx_3^2}{2}, \quad \chi_v = e^{\frac{i\pi}{4}} T_v. \quad (26)$$

Similarly, we find

$$\begin{aligned} {}_+\psi_{rp_v}(x|A_3 = -Et) &= \int_{-\infty}^{\infty} dp^- N(p_v, p^-) {}_+\psi_{rp^-}(x|A_3 = -Et) = \\ &= \frac{\exp[-\frac{\pi\varkappa}{4}]}{2\sqrt{eE\varkappa(1-v)}} [u_r D_{i\varkappa}(-\chi_v^*) + e^{-\frac{i\pi}{4}} \sqrt{\frac{2eE(1-v)}{1+v}} \varkappa u'_r D_{i\varkappa-1}(-\chi_v^*)] \times \\ &\exp\{i[\frac{\pi}{2} + \frac{\varkappa}{2} \ln \frac{1+v}{2(1-v)} + \tilde{\vartheta}_v]\}, \quad \chi_v^* = e^{-i\frac{\pi}{4}} T_v. \end{aligned} \quad (27)$$

We note that ${}_-\psi_{p_v}$ (${}_-\psi_{p_v}$) can be obtained from ${}_+\psi_{p_v}$ (${}_+\psi_{p_v}$) by substitutions $T_v \rightarrow -T_v$, $u'_r \rightarrow -u'_r$.

To compare (26) with (14-15), we go in (26) to the spinor representation. From (18) and (20) we have

$$Uu'_1 = \sqrt{2} \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \quad Uu'_2 = \sqrt{2} \begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \end{bmatrix}, \quad Uu_1 = \sqrt{2} \begin{bmatrix} p_1 - ip_2 \\ 0 \\ 0 \\ m \end{bmatrix}, \quad Uu_2 = \sqrt{2} \begin{bmatrix} m \\ 0 \\ 0 \\ -p_1 - ip_2 \end{bmatrix}. \quad (28)$$

Now we see that (26) indeed agrees with (14).

Thus, the transformation (24) relates the solutions ψ_{rp^-} to ψ_{p_v} . For $v = 0$ the latter solutions become solutions ψ_{rp_3} and they coincide up to an unimportant phase factor $\exp[i\frac{\pi}{2} - \frac{i\varkappa}{2} \ln 2]$ with the normalized solutions obtained in [9], see also [4]. This justifies the normalization factors and the classification of ψ_{rp^-} solutions in (19).

We see that the classification in ψ_{rp_3} (which is beyond doubt) determines the classification in ψ_{rp_v} and ψ_{rp^-} . In the next Section we shall see that the classification in ψ_{rp^-} in its turn determines the classification in ψ_{rp_s} and ψ_{rp^0} .

3 Stationary solutions

These solutions appear naturally when we use instead of the vector-potential (2) the potential

$$A^0 = -Ex_s, \quad x_s = x_3 - st, \quad 0 \leq s < 1. \quad (29)$$

The solutions of the squared Dirac equation (3) can be obtained from the corresponding solutions of the Klein-Gordon equation (see eq. (59) in [6]) by the same substitutions (7). For positive-frequency in-solution we find

$${}_+F = \exp[i\vartheta_s] \text{diag}(\varphi_1, \varphi_2, \varphi_2, \varphi_1), \quad \varphi_1 = D_{i\varkappa}(\chi_s), \quad \varphi_2 = D_{i\varkappa-1}(\chi_s), \quad \chi_s = e^{\frac{i\pi}{4}} Z_s, \quad (30)$$

$$Z_s = \sqrt{\frac{2}{eE(1-s^2)}} (eEx_s - p_s), \quad p_s = p^0 - sp_3, \quad \vartheta_s = -\frac{p_s(t - sx_3)}{1-s^2} - \frac{eEsx_s^2}{2(1-s^2)}. \quad (31)$$

The solutions of the Dirac equation are given by the columns of the matrix in eq. (10), where now

$$\Pi^0 = i \frac{\partial}{\partial t} - eEx_s, \quad \Pi_3 = -i \frac{\partial}{\partial x_3}.$$

From here and (30) we get

$$(\Pi^0 - \Pi_3)e^{i\vartheta_s}\varphi(\chi_s) = -e^{i\vartheta_s - \frac{i\pi}{4}} \sqrt{\frac{2eE(1-s)}{1+s}} \left(\frac{\partial}{\partial \chi_s} + \frac{\chi_s}{2} \right) \varphi(\chi_s). \quad (32)$$

Using again the fourth and the thirist columns in (9), we obtain in the spinor representation

$${}^+\psi_{1p_s}(x|A^0 = -Ex_s) \propto \begin{bmatrix} (p_1 - ip_2)D_{i\kappa}(\chi_s) \\ -e^{\frac{i\pi}{4}} \sqrt{\frac{2eE(1-s)}{1+s}} \kappa D_{i\kappa-1}(\chi_s) \\ 0 \\ mD_{i\kappa}(\chi_s) \end{bmatrix} e^{i\vartheta_s}, \quad (33)$$

$${}^+\psi_{2p_s}(x|A^0 = -Ex_s) \propto \begin{bmatrix} mD_{i\kappa}(\chi_s) \\ 0 \\ -e^{\frac{i\pi}{4}} \sqrt{\frac{2eE(1-s)}{1+s}} \kappa D_{i\kappa-1}(\chi_s) \\ -(p_1 + p_2)D_{i\kappa}(\chi_s) \end{bmatrix} e^{i\vartheta_s}. \quad (34)$$

The normalized solutions ψ_{rp_s} can be obtained from ψ_{rp^-} in (19) by the same integral transformation as in the scalar case [6]

$$\psi_{rp_s} = \int_{-\infty}^{\infty} dp^- S(p_s, p^-) \psi_{rp^-}, \quad (35)$$

$$S(p_s, p^-) = (2\pi eE(1-s))^{-\frac{1}{2}} \exp\left\{-i \frac{[p^-(1+s)]^2 - 4p^-p_s(1+s) + 2p_s^2}{4eE(1-s^2)}\right\}, \quad (36)$$

For the potential

$$\mathcal{A}_\mu = \delta_{\mu 0} Ex_3, \quad \mathcal{A}^0 = -\mathcal{A}_0. \quad (37)$$

we find in standard representation

$${}^+\psi_{rp_s}(x|\mathcal{A}) = \frac{\exp[-\frac{3\pi\kappa}{4}]}{2\sqrt{eE(1-s)}} [u_r D_{-i\kappa-1}(\chi_s^*) - e^{\frac{i\pi}{4}} \sqrt{\frac{2eE(1-s)}{1+s}} u'_r D_{-i\kappa}(\chi_s^*)] \times \\ \exp\left\{i\left[\frac{\pi}{2} + \frac{\kappa}{2} \ln \frac{1+s}{2(1-s)} + \tilde{\vartheta}_s\right]\right\}, \quad \tilde{\vartheta}_s = \vartheta_s - \frac{eEst^2}{2}, \quad \chi_s^* = e^{-\frac{i\pi}{4}} Z_s, \quad (38)$$

$${}^+\psi_{rp_s}(x|\mathcal{A}) = \frac{\exp[-\frac{3\pi\kappa}{4}]}{2\sqrt{eE\kappa(1-s)}} [u_r D_{i\kappa}(\chi_s) - e^{\frac{i\pi}{4}} \sqrt{\frac{2eE(1-s)}{1+s}} \kappa u'_r D_{i\kappa-1}(\chi_s)] \times \\ \exp\left\{i\left[\frac{\pi}{2} + \frac{\kappa}{2} \ln \frac{1+s}{2(1-s)} + \tilde{\vartheta}_s\right]\right\}. \quad (39)$$

The solutions ${}^-\psi_{rp_s}$ (${}^-\psi_{rp_s}$) can be obtained from ${}^+\psi_{rp_s}$ (${}^+\psi_{rp_s}$) by substitutions $Z_s \rightarrow -Z_s$, $u'_r \rightarrow -u'_r$. The solutions ψ_{rp_s} satisfy the same Bogoliubov transformations (23) as ψ_{rp_v} .

For $s = 0$ we get the usual stationary solutions with the quantum number p^0 . Their normalization factor can be checked independently (up to an inessential phase factor) using the relation between the normalization on unit current and the usual normalization. This relation for the scalar particle was established in [6]. The result remains valid (up to a sign for negative-frequency states) for the spinor particle as well. To apply this result, we first relate our classification to the classification according to the sign of momentum in asymptotic regions $x_3 \rightarrow \pm\infty$. (In this classification the sign of particle momentum along x_3 plays the role of sign of energy [4].) To distinguish this classification from the previous one, we denote the solutions in this classification as f_n , (n is the set of quantum numbers) and assume that they are normalized on a unit current (up to a sign) along x_3 . The correspondence between functions in this two classifications is given by

$${}^-\psi_n \propto {}_+f_n, \quad {}^-\psi_n \propto -f_n, \quad {}^+\psi_n \propto {}_+f_n, \quad {}^+\psi_n \propto -f_n. \quad (40)$$

Here ${}_+f_n$ ($-f_n$) is the solution with only one wave with the positive (negative) momentum along x_3 for $x_3 \rightarrow -\infty$. Similarly, for $x_3 \rightarrow \infty$ the sign of momentum is indicated in the upper position.

In terms of f_n eq.(23) can be rewritten as

$$\begin{aligned} {}_+f_n &= c'_{1n} {}^+f_n + c'_{2n} {}^-f_n, \quad {}^+f_n = -c'^*_{1n} {}_+f_n - c'_{2n} {}^-f_n; \\ |c'_{2n}|^2 - |c'_{1n}|^2 &= 1, \quad c'_{1n} = -\frac{c_{1n}}{c_{2n}}, \quad c'_{2n} = \frac{1}{c_{2n}^*} \end{aligned} \quad (41)$$

(see eqs. (5.17), (5.18) in [4]; ψ_n there are f_n here). Now the above-mentioned relation between usual normalization and normalization on the unit current can be written in the form

$$\int dx_3 J^0({}_+f_{rp^0}, {}_+f_{rp^0}) = |c'_{2p}|^2 2\pi \delta(p'^0 - p^0). \quad (42)$$

So, to obtain the normalization on $2\pi\delta(p'^0 - p^0)$, we have to divide the solution f_n by $|c'_{2n}|$. To normalize f_n on unit current, we use as f_n the expression (33) or (34) in

$$J_3 = f_n^* \alpha_3 f_n. \quad (43)$$

Then we use the relation

$$D_{-i\kappa}(y) D_{i\kappa}(iy) - \kappa D_{i\kappa-1}(iy) D_{-i\kappa-1}(y) = e^{-\frac{\pi\kappa}{2}}, \quad (44)$$

which follows from (see eq. (8.2.11) in [7])

$$D_\nu(y) \frac{d}{dy} D_{-\nu-1}(iy) - D_{-\nu-1}(iy) \frac{d}{dy} D_\nu(y) = -ie^{-i\frac{\pi\nu}{2}} \quad (45)$$

and relations (13). In this way we find in the spinor representation, where α_3 is given in (4), the normalization factor

$$|N_{rp^0}| = \frac{e^{\frac{\pi\kappa}{4}}}{\sqrt{m^2 + p_1^2 + p_2^2}} \frac{1}{|c'_{2n}|} = \frac{e^{-\frac{3\pi\kappa}{4}}}{\sqrt{2eE\kappa}}. \quad (46)$$

This is in agreement with (39) for $s = 0$ (and with the expression for ${}^-\psi_{p^0}$ obtainable from ${}^+\psi_{p^0}$ as indicated below eq.(39)), if we go there from standard to spinor representation with the help of (28).

4 Concluding remarks

In this paper we have completed the study of equivalent sets of solutions of the Klein-Gordon and Dirac equations with constant electric field initiated in [10, 11]. We find that in the spinor case everything is just as expected from the scalar case, considered in [6].

5 Acknowledgments

This work was supported in part by the Russian Foundation for Basic Research (projects no 00-15-96566 and 02-02-16944).

References

1. A.Hansen, F.Ravndal, *Physica Scripta*, **23**, 1036 (1981).
2. W.Greiner, B.Müller, J.Rafelski, *Quantum Electrodynamics of Strong Field*, Springer-Verlag (1985).
3. A.Calogeracos, N.Dombey, *Contemp. Phys.* **40**, 313 (1999).
4. A.I.Nikishov, *Tr. Fiz. Inst. Akad. Nauk SSSR* **111**, 152 (1979) ;
J. Sov. Laser Res. **6**, 619 (1985).
5. A.I.Nikishov, hep-th/0111137; *Problems of Atomic Science and Technology*, 2001, N6, p. 103.
6. A.I.Nikishov, hep-th/0202024
7. *Higher Transcendental Functions, Vol. 2 (Bateman Manuscript Project)*,
Ed. by A.Erdélyi (McGraw-Hill, New York, 1953; Nauka, Moscow, 1980; Pergamon, Oxford, 1982).
8. L.H.Ryder, *Quantum Field Theory* (Cambridge University Press, 1985)
9. A.I.Nikishov, *Zh. Eksp. Teor. Fiz.* **57**, 1210 (1969); *Sov. Phys.-JETP* **30**, 660 (1970).
10. N.B.Narozhny and A.I.Nikishov, *Teor. Mat. Fiz.* **26**, 16 (1976).

11. N.B.Narozhny and A.I.Nikishov, Tr. Fiz. Inst. Akad. Nauk SSSR **168**, 175 (1985);
in *Issues in Intensive-Field Quantum Electrodynamics*, Ed. by V.L.Ginzburg (Nova
Science, Commack, 1987).